

On the wave equation in spacetimes of Gödel type

P Marecki

ITP, Universitaet Leipzig, Postfach 100 920, D-04009 Leipzig, Germany

E-mail: pmarecki@gmail.com

Abstract. We analyze the d'Alembert equation in the Gödel-type spacetimes with spherical and Lobachevsky sections (with sufficiently rapid rotation). By separating the t and x_3 dependence we reduce the problem to a group-theoretical one. In the spherical case solutions have discrete frequencies, and involve spin-weighted spherical harmonics. In the Lobachevsky case we give simple formulas for obtaining all the solutions belonging to the $D_\lambda^\pm$ sectors of the irreducible unitary representations of the reduced Lorentz group. The wave equation enforces restrictions on λ and the allowed (here: continuous) spectrum of frequencies.

1. Introduction

When a rapidly rotating gas of bosons is cooled down below the Bose-Einstein condensation temperature a regular lattice of vortices (Abrikosov lattice) emerges. The velocity vector field which initially circulated around the only axis of rotation now rotates locally around every vortex core. Interestingly, in the General Relativity Theory there also exist both types of rotation. A homogeneous spacetime with the latter type of rotation was discovered by Gödel [1] long time ago. Chapline and Mazur have argued recently [2] that a BEC with an array of vortices and a Gödel-type metric will provide a solution of the Einstein-Gross-Pitaevskii system (in the coarse-grained, $v \ll c_s$ limit). By Gödel-type we mean homogeneous stationary spacetimes in which matter rotates around every point relative to the “compass of inertia” [3, 4]. As in isotropic cosmology, the hypersurfaces “ $t = \text{const}$ ” can have either Lobachevsky, flat or spherical geometry[‡] with the radius of curvature being a parameter. The angular velocity of rotation provides the second parameter of the Gödel-type spacetimes.

In this paper we set ourselves a modest task of analyzing the d'Alembert equation in the Gödel-type spacetimes with spherical and Lobachevsky sections (in the latter case we confine ourselves to the case of sufficiently rapid rotation). We shall, of course, profit from the work of a number of authors who have, from different angles, attacked this problem before, [5, 6, 7, 8, 9, 10]. Therefore, our goal here is not to discover, but rather to consolidate, simplify (and where necessary - to correct) the work of others in order to lay solid foundations for further progress.

This short paper has one main section (the second one), which is divided into two parts treating the d'Alembert equation in the Lobachevsky and spherical cases separately. We have also included appendices on the Lobachevsky geometry and on the sources of Gödel-type spacetimes for quick reference.

[‡] The type of geometry is controlled by, among other things, the sign of the cosmological constant. The spherical one corresponds to a positive vacuum energy currently supported on the cosmological scale by multiple observations.

2. Solving the wave equation by an Ansatz

In order to find the solutions of the d'Alembert equation in the spacetimes of Gödel type we will make use of the fact, that these spacetimes are stationary, homogeneous and axially symmetric. Thus, we will assume that the solutions, Ψ , of the wave equation, $\square\Psi = 0$, can be separated according to

$$\Psi(t, x, \varphi, x_3) = e^{-i\omega t} e^{ik_3 x_3} \cdot \psi(x, \varphi), \quad (1)$$

where the function $\psi(x, \varphi)$ should be periodic in φ with the period of 2π . In the following we shall classify the allowed functions, $\psi(x, \varphi)$, in the Lobachevsky and spherical cases separately.

2.1. Lobachevsky case

The original Gödel spacetime is a stationary four-dimensional spacetime with the $t = \text{const}$ hypersurfaces being just $\mathbb{H}_2 \times \mathbb{R}$, where $\mathbb{H}_2$ is the Lobachevsky space (or hyperbolic plane, see appendix A). These hypersurfaces are threaded by a constant, homogeneous gravimagnetic field perpendicular to the Lobachevsky space (i.e. along the x_3 direction). There family of Gödel-type metrics [1] is parameterized by the “amount of rotation” Ω and the magnitude of Lobachevsky curvature denoted by l (see appendix B):

$$d(2ls)^2 = (dt + Hd\varphi)^2 - dx^2 - D^2 d\varphi^2 - dx_3^2, \quad (2)$$

where all the coordinates are dimensionless, and

$$H = 2\alpha \sinh^2(x/2) \quad (3)$$

$$D = \sinh(x), \quad (4)$$

with $\alpha = \Omega/l$. Both t and x_3 range over the whole $\mathbb{R}$, while $x \in [0, \infty)$ and $\varphi \in [0, 2\pi)$ with the usual identification of 0 and 2π . The metric is of the form $d\tau^2 = (dt + A_i dx^i)^2 - h_{ij} dx^i dx^j$, with the Lobachevsky space metric h_{ij} and a constant gravimagnetic (Lense-Thirring) field on this space

$$B^i = \frac{\epsilon^{ijk} \partial_j A_k}{\sqrt{h}} = 2\alpha, \quad (5)$$

with the same constant being equal to the contravariant x_3 component of the rotation vector of the $(\partial_t)^\alpha$ vector field (velocity field of the sources). The case $\Omega = 0$ corresponds to a static spacetime with Lobachevsky spatial sections, while $\Omega^2 = 2l^2$ corresponds to the original Gödel metric.

In this paper we deal only with spacetimes of sufficiently rapid rotation $\Omega > l$ (that is, $\alpha > 1$). In order to solve the wave equation in these spacetimes, we will make use of the underlying Lobachevsky structure. The d'Alembert equation for Ψ is

$$\begin{aligned} \square\Psi = [1 - \alpha^2 \tanh^2(x/2)] \Psi_{tt} + \frac{\alpha}{\cosh^2(x/2)} \Psi_{t\varphi} - \\ - \frac{1}{\sinh^2(x)} \Psi_{\varphi\varphi} - \Psi_{33} - \Psi_{xx} - \coth(x) \Psi_x = 0. \end{aligned} \quad (6)$$

The following three operators, which are Killing vectors of the Gödel spacetimes multiplied by $-i$, fulfill the same commutation relations as the Killing vectors L_i

of the Lobachevsky plane (see appendix A):

$$J_1 = -i \sin \varphi \partial_x - i \coth(x) \cos(\varphi) \partial_\varphi - i \alpha \cos \varphi \tanh(x/2) \partial_t, \quad (7)$$

$$J_2 = i \cos \varphi \partial_x - i \coth(x) \sin(\varphi) \partial_\varphi - i \alpha \sin \varphi \tanh(x/2) \partial_t, \quad (8)$$

$$J_0 = -i \partial_\varphi - i \alpha \partial_t, \quad (9)$$

while the remaining two Killing vectors, ∂_t and ∂_3 , obviously commute with the above ones. We notice further, that making use of the separation ansatz (1) for Ψ , we are lead to

$$\square \Psi = -C_2 \Psi - (1 - \alpha^2) \omega^2 \Psi - \Psi_{33}, \quad (10)$$

where here: $C_2 = J_0^2 - J_1^2 - J_2^2$. The wave equation therefore reduces remarkably to

$$C_2 \psi(x, \varphi) = [(\alpha^2 - 1) \omega^2 + k_3^2] \psi(x, \varphi) \quad (11)$$

Because we are interested in the regime $\alpha^2 > 1$, it remains to determine the possible eigenfunctions, $\psi(x, \varphi)$, with a *positive* eigenvalue of the Casimir operator C_2 which was defined above. Because of the ansatz, the operator C_2 becomes now a second order partial differential operator acting on $\psi(x, \varphi)$.

In order to determine the eigenfunctions, let us now regard C_2 (and, in fact, the whole set J_a) as an operator acting in the Hilbert space of square-integrable functions of x, φ with the scalar product provided by the Lobachevsky product, (A.4).

It is helpful to recall the classification of the unitary irreducible representations of the reduced Lorentz group $\mathfrak{L}_3$ (two boosts and one rotation) and of its corresponding spinor group $\mathfrak{S}_3 \equiv \text{SU}(1,1)$ of Bargmann [11]. For positive eigenvalues of C_2 one is lead to what Bargmann calls *discrete classes*§, D_λ^- and D_λ^+ . Within $D_\lambda^\pm$ the eigenvalue of C_2 is $\lambda(\lambda \mp 1)$ and the eigenvalues of J_0 start with λ and increase (without bound) by one with each application of $J_+ = J_1 + iJ_2$ in the D_λ^+ class (or, respectively, start with $-\lambda$ and decrease (without bound) by one with each application of J_- , for the D_λ^- class). In D_λ^+ the number λ is real and positive||, while in D_λ^- it is real and negative.

Let us determine all these functions explicitly. For the complex parameter

$$z = \tanh(x/2) e^{i\varphi}, \quad (12)$$

we note that

$$\partial_\varphi = iz \partial - i \bar{z} \bar{\partial},$$

where $\bar{\partial} = \partial_{\bar{z}}$. We find

$$J_+ = z^2 \partial - \bar{\partial} + az, \quad (13)$$

$$J_- = \partial - \bar{z}^2 \bar{\partial} + a \bar{z}, \quad (14)$$

$$J_0 = z \partial - \bar{z} \bar{\partial} + a, \quad (15)$$

where

$$a = -\alpha \omega \quad (16)$$

and $J_\pm = J_1 \pm iJ_2$. We may verify, that

$$[J_+, J_-] = -2J_0, \quad (17)$$

$$C_2 = J_0(J_0 + 1) - J_- J_+, \quad (18)$$

$$C_2 = J_0(J_0 - 1) - J_+ J_-, \quad (19)$$

§ Note, that our convention on the sign of the Casimir operator, C_2 , is different from that of [11].

|| Bargmann notes (see §4 of [11]), that the universal covering group $\mathfrak{C}$ covers $\mathfrak{S}_3$ infinitely often. He restricts himself to single-valued representations of $\mathfrak{S}_3$, and as a consequence treats only integer and half-integer λ 's. We see no reason for such a restriction here, and therefore leave λ real.

which are all standard relations in the theory of representations of $SU(1,1)$.

For a given λ the families of simultaneous eigenfunctions of C_2 and J_0 are constructed upon a reference vectors, denoted here by ψ_0^+ (or ψ_0^-), annihilated by $\P$ (J_- (or J_+)). This leads to two families, D_λ^+ and D_λ^- .

2.1.1. Structure of D_λ^+ . We find immediately

$$\psi_0^+ = (1 - z\bar{z})^\lambda \bar{z}^{a-\lambda}, \quad (20)$$

where λ is the eigenvalue of J_0 , and $\lambda(\lambda - 1)$ the eigenvalue of C_2 . The function ψ_0^+ depends, via the last term, on the angle φ . Since it must be periodic in φ , with identical values for $\varphi = 0$ and $\varphi = 2\pi$, the number $a - \lambda \equiv m$ must be integer. In the original variables we have

$$\psi_0^+ = \cosh(x/2)^{-2\lambda-m} \sinh(x/2)^m e^{-im\varphi}. \quad (21)$$

Further restrictions,

$$m \geq 0, \quad \lambda > \frac{1}{2}, \quad (22)$$

are obtained from the requirement of square integrability w.r.t. the product (A.4) at $x = 0$ and at infinity, respectively⁺.

A connection with the methods of ordinary differential equations is provided by the following consideration: the eigenvalue problem for the Casimir operator,

$$C_2\psi = \lambda(\lambda - 1)\psi, \quad (23)$$

with $\lambda \geq 1$ leads, with a separation ansatz $\psi = f(x) \exp(-im\varphi)$, $\mathbb{Z} \ni m \geq 0$ to the equation

$$w(w-2)f'' + 2(w-1)f' + \left[\frac{2a(a+m)}{w} - \frac{m^2}{w(w-2)} \right] f = \lambda(\lambda-1)f, \quad (24)$$

where $w = 2 \cosh^2(x/2)$. This is an equation with three regular singular points, at $(0, 2, \infty)$, with the exponents $(a + \frac{m}{2}, a - \frac{m}{2})$, $(\frac{m}{2}, -\frac{m}{2})$ and $(\lambda - 1, -\lambda)$ respectively. Evidently, the function(s) we have found by algebraic methods, ψ_0^+ , have the right asymptotics at $w = 2$ ($\psi_0^+ \sim (w-2)^{\frac{m}{2}}$) and at infinity ($\psi_0^+ \sim w^{-\lambda}$) to be square integrable at these extremes.

Let us now summarize the results. The solutions are characterized by the momentum in the x_3 direction (k_3) and by the representation parameters (λ, k) with $k \in \{\lambda, \lambda + 1, \dots, \infty\}$. The wave equation has the form

$$\lambda(\lambda - 1) = \beta^2 a^2 + k_3^2, \quad (25)$$

where $\beta^2 = 1 - 1/\alpha^2 > 0$ depends only on the parameter α of the spacetime, and $a - \lambda \geq 0$ is an integer. The above dispersion relation is unusual by the sign in front of k_3^2 . The solution for a is given by

$$a = \frac{1}{\beta} \sqrt{\lambda(\lambda - 1) - k_3^2}, \quad (26)$$

but only if $a \geq \lambda$, which is possible only if $\lambda \geq \alpha^2$. Consequently, because of (16), there also exists a gap in frequencies: $|\omega| \geq \alpha$.

[¶] The necessity for such a vector is clear; as in the case of the rotation group - its absence would lead to negative-norm states created by successive application of either J_- (or J_+).

⁺ Even though on the surfaces of constant t and x_3 the ∂_φ vector becomes time-like for large x , allowing for functions growing exponentially with x (omitting the second restriction) would certainly make no sense.

2.1.2. *Structure of D_λ^- .* The function

$$\psi_0^- = (1 - z\bar{z})^{-\lambda} z^{\lambda-a}, \quad (27)$$

is an eigenfunction of C_2 to the eigenvalue $\lambda(\lambda+1)$, and of J_0 to the eigenvalue λ . It is square-integrable if $\lambda < -1/2$ and if $m = \lambda - a$ (which is an integer) is non-negative. Consequently, C_2 is non-negative and ψ_0^- is square-integrable if $a \leq \lambda \leq -1$. Because of that, the frequency ω is now positive, and the wave equation

$$\lambda(\lambda+1) = \beta^2 a^2 + k_3^2, \quad (28)$$

has solutions

$$a = -\frac{1}{\beta} \sqrt{\lambda(\lambda+1) - k_3^2}, \quad (29)$$

only if $\lambda \leq -\alpha^2$ (in order to satisfy $a \leq \lambda$). Here, again $|\omega| \geq \alpha$.

2.2. Spherical case

The case of spacetimes with spherical (x, φ) hyperplanes (threaded by a homogeneous gravimagnetic field) is simpler, and the resulting functions, $\psi(x, \varphi)$ belong to a known family of functions (spin “a” spherical harmonics). In this case, $H = 2\alpha \sin^2(x/2)$ and $D = \sin(x)$ (where $x \in [0, \pi)$ has the interpretation of the latitude). The wave equation can be solved similarly, though in this case the d'Alembert operator is given by

$$\square \Psi = [C_2 - (1 + \alpha^2)\omega^2 + k_3^2] \Psi = 0, \quad (30)$$

where C_2 is the Casimir operator of a modification of the algebra the $SO(3)$ of (symmetry) generators:

$$L_+ = z^2 \partial + \bar{\partial} + az, \quad (31)$$

$$L_- = -\partial - \bar{z}^2 \bar{\partial} + a\bar{z}, \quad (32)$$

$$L_3 = z\partial - \bar{z}\bar{\partial} + a, \quad (33)$$

and

$$z = \tan(x/2)e^{i\varphi}.$$

The above operators fulfill the $SO(3)$ relations:

$$[L_+, L_-] = 2L_3, \quad [L_\pm, L_3] = \mp L_\pm.$$

The representation spaces are grouped in sectors, each with a definite value of $L^2 = \lambda(\lambda+1)$ and $L_3 \in [-\lambda, \lambda]$, where λ is a positive integer or half-integer*.

The lowest states, $\psi_{-\lambda}$, which are annihilated by L_- with the eigenvalue $-\lambda$ of L_3 , and the highest states ψ_λ (annihilated by L_+ with the eigenvalue λ of L_3) can now be determined. One finds

$$\psi_{-\lambda} = (1 + z\bar{z})^{-\lambda} \bar{z}^{\lambda+a}, \quad (34)$$

$$\psi_\lambda = (1 + z\bar{z})^{-\lambda} z^{\lambda-a}. \quad (35)$$

* Only the integer or half integer values of λ are allowed, for otherwise one would be lead to standard contradictions (eg. states of negative norm, after sufficiently many applications of L_+).

These function must be periodic in φ , with identical values for $\varphi = 0$ and $\varphi = 2\pi$, and therefore $a + \lambda \equiv m$ must also be an integer. Now it is easy to see, that either there holds

$$m \geq 0, \quad (36)$$

$$m \leq 2\lambda, \quad (37)$$

or the function $\psi_{-\lambda}$ will not be square integrable on the sphere (in x, φ) because of singularities at $x = 0$ or $x = \pi$ respectively. Thus, for a given value of $\lambda \in \{\frac{1}{2}, 1, \dots, \infty\}$ there remains a freedom to choose $m \in [0, 2\lambda]$. This can be viewed as a condition specifying a , so that the structure is the following:

There are infinitely many sectors with integer or half-integer a . In each of these sectors there are eigenspaces of L^2 to the eigenvalues $\lambda(\lambda + 1)$ with $\lambda \geq |a|$ (with an integer $\lambda + a$). Each of such eigenspaces is spanned by $2\lambda + 1$ functions generated by a successive application of L_+ to $\psi_{-\lambda}$.

The eigenspace $\lambda = \frac{1}{2}$, $a = -\frac{1}{2}$ consists of two functions first encountered by Dirac [12] in the context of magnetic monopoles,

$$\psi_0 = \cos(x/2) \quad \text{for } k = -\frac{1}{2}, \quad (38)$$

$$\psi_1 = -\sin(x/2)e^{i\varphi} \quad \text{for } k = \frac{1}{2}. \quad (39)$$

The functions we have found are also known as the monopole harmonics or the spin- a spherical harmonics [13, 14], so that further properties (eg. completeness for each a) and identities among these functions are known.

We may now summarize the results taking into account the wave equation. Each solution is characterized by the parameters: frequency (ω), momentum in the x_3 direction (k_3), sector number a (which is an integer or half-integer), and the pair (λ, k) with $\lambda \geq |a|$ and $k \in [-\lambda, \lambda]$. With

$$\omega = \frac{a}{\alpha} \quad (40)$$

the frequency is fixed by the sector number a . Such a quantization of frequencies has an alternative, kinematical (gauge) derivation pointed out long time ago by Mazur [15]. The only role of the wave equation is to provide a relation between k_3 , λ and a :

$$\lambda(\lambda + 1) + k_3^2 = (1 + 1/\alpha^2)a^2. \quad (41)$$

Thus, in each sector (for each energy) k_3 belongs to a discrete bounded set.

3. Conclusions

Starting with the separation ansatz (1) we have solved the d'Alembert equation in the spacetimes of Gödel type. In the spherical case the functions with the frequency $\omega = n/2\alpha$, $n \in \mathbb{Z}$ turned out to coincide with the $n/2$ -weighted spherical harmonics. In the Lobachevsky case we have given simple formulas for obtaining all the solutions belonging to the $D_\lambda^\pm$ sectors of irreducible unitary representations of the reduced Lorentz group. The wave equation enforced $\alpha^2 \leq |\lambda| \in \mathbb{R}$ and a gap in frequencies $\omega \in \mathbb{R} \setminus (-\alpha, \alpha)$.

Acknowledgements

I would like to thank Rainer Verch and Jan Schlemmer for several useful suggestions.

References

- [1] Gödel K, 1949 *Rev. Mod. Phys.* **21**, 447
- [2] Chapline, G and Mazur P O, 2004 *gr-qc/0407033*
- [3] Kundt W, 1956 *Z. Phys.* **C145**, 611
- [4] Kajari E, Walser R, Schleich W P, Delgado A, 2004 *Gen. Relat. Gravit.* **36**, 2289
- [5] Mashoon B, 1975 *Phys. Rev.* **D11**, 2679
- [6] Hiscock W, 1978 *Phys. Rev.* **D17**, 1497
- [7] Comtet A, 1987 *Ann. Phys. (NY)* **173**, 185
- [8] Dunne G V, 1992 *Ann. Phys. (NY)* **215**, 233
- [9] Figueiredo B, Soares D and Tiomno J, 1992 *Class. Quant. Grav.* **9**, 1593
- [10] Dukker N, Fiol B and Simón J, 2004 *JCAP* **10**, 12
- [11] Bargmann V, 1947 *Ann. Math.* **48**, 568
- [12] Dirac P A M, 1931 *Proc. Roy. Soc. Lond. A* **133**, 60
- [13] Tamm Ig, 1931 *Zeitschrift fuer Physik A* **71**, 141
- [14] Dray T 1984 *J. Math. Phys.* **26**, 1030
- [15] Mazur P O 1986 *Phys. Rev. Lett.* **57**, 929
- [16] Raychaudhuri A K, and Guha Thakurta S N 1980 *Phys. Rev. D* **22**, 802
- [17] Rebouças M J, and Tiomno J 1983 *Phys. Rev. D* **28**, 1251

Appendix A. Preliminaries on the Lobachevsky space

The 2-dimensional Lobachevsky space (known also as the hyperbolic plane, $\mathbb{H}_2$) is a space defined as the hypersurface $t^2 - x^2 - y^2$ of the 1+2 dimensional Minkowski space. The intrinsic metric of this space is given by

$$ds^2 = dr^2 + \sinh^2(r) d\varphi^2,$$

with $r \in [0, \infty)$, $\varphi \in [0, 2\pi)$. (This map covers the whole space.) The three Killing vector fields,

$$K_1 = \sin(\varphi)\partial_r + \coth(\psi)\cos(\varphi)\partial_\varphi \quad (\text{A.1})$$

$$K_2 = -\cos(\varphi)\partial_r + \coth(\psi)\sin(\varphi)\partial_\varphi \quad (\text{A.2})$$

$$K_0 = \partial_\varphi, \quad (\text{A.3})$$

by complete analogy with rotations on a sphere, when multiplied by $-i$ lead to the generators of the respective symmetries on the Hilbert space of square integrable functions on $\mathbb{H}_2$, with the standard product

$$(f, g) = \int \bar{f}g \sinh(r) dr d\varphi. \quad (\text{A.4})$$

These generators, denoted here by L_i , fulfill the commutation relations of the algebra of $\text{SU}(1,1)$ generators:

$$[L_0, L_1] = iL_2 \quad (\text{A.5})$$

$$[L_2, L_0] = iL_1 \quad (\text{A.6})$$

$$[L_1, L_2] = -iL_0, \quad (\text{A.7})$$

and commute with the quadratic Casimir operator

$$C_2 = L_0^2 - L_1^2 - L_2^2. \quad (\text{A.8})$$

The group $\text{SU}(1,1)$ is not compact, and therefore the representation theory is richer than that of the rotation group. Depending on the sign of the eigenvalue of C_2 there are, discrete or continuous series of eigenfunctions. Let us finally notice, that

$$C_2 = \partial_r^2 + \coth(r)\partial_r + \frac{1}{\sinh^2(r)}\partial_\varphi^2 = \frac{1}{\sqrt{h}}\partial_i \left(h^{ij} \sqrt{h} \partial_j \right) \quad (\text{A.9})$$

is just the Laplace-Beltrami operator on the Lobachevsky space, and therefore here $C_2 \leq 0$.

Appendix B. Sources

Appendix B.1. Spherical case

Let us choose the orthonormal tetrad

$$\begin{aligned} \mathbf{e}^0 &= dt + m d\varphi, \\ \mathbf{e}^\Phi &= D d\varphi, \\ \mathbf{e}^R &= dr, \\ \mathbf{e}^3 &= dx^3, \end{aligned}$$

where the functions m and D are as in section 2.2. The spherical Gödel spacetime can be generated by [16, 17] a (neutral) dust, supplemented by a cosmological constant (with a positive vacuum energy) and an (source-free) electromagnetic field directed along the x_3 axis:

$$\begin{aligned} F^{R\Phi} &= E \cos(\alpha x_3), \\ F^{30} &= E \sin(\alpha x_3). \end{aligned}$$

The Einstein field equations

$$G_{AB} = \kappa(\rho u_A u_B + T_{AB}) - \Lambda \eta_{AB} \quad (\text{B.1})$$

with $\kappa = 8\pi G/c^4$, and the dust four-velocities, u_A , coincident with the $(\partial_t)^a$ (and with e_0^a) vector fields. We get

$$\begin{aligned} G_{AB} &= \text{diag}(1 + 3A, A, A, -1 - A), \\ T_{AB} &= \frac{E^2}{8\pi} \text{diag}(1, 1, 1, -1), \\ -\Lambda \eta_{AB} &= -\Lambda \text{diag}(1, -1, -1, -1), \end{aligned}$$

where $A = \alpha^2/4$. After an additional rescaling of the line element $ds^2 \rightarrow R^2 ds^2$, with $R = 1/2l$ being the radius of the sphere, we find

$$\begin{aligned} \Lambda &= -\frac{1}{2R^2}, \\ \rho &= \frac{\alpha^2 c^4}{16\pi G R^2}, \\ E^2 &= \frac{c^4}{4G R^2} (2 + \alpha^2). \end{aligned}$$

Alternatively, by introducing the vacuum energy, $\rho_0 = -\Lambda c^4/8\pi G > 0$, we may write the last two equations as

$$\begin{aligned} \rho &= \alpha^2 \rho_0, \\ E^2 &= 4\pi \rho_0 (2 + \alpha^2). \end{aligned}$$

Thus, in the spherical Gödel-type spacetime the vacuum energy controls the scale R , the dust's density the rotation, while the energy density of the electromagnetic field needs to be of the order of the vacuum energy.

Appendix B.2. Lobachevsky case

We use the same tetrad form, but now with the functions m and D from section 2.1. We get

$$G_{AB} = \text{diag}(-1 + 3A, A, A, 1 - A),$$

$$T_{AB} = \frac{E^2}{8\pi} \text{diag}(1, 1, 1, -1),$$

and the Einstein equations lead to

$$\Lambda = \frac{1}{2R^2},$$

$$\rho = \frac{\alpha^2 c^4}{16\pi G R^2},$$

$$E^2 = \frac{c^4}{4GR^2}(\alpha^2 - 2).$$

For $\alpha^2 = 2$ we obtain the Gödel's original solution, which is generated by the dust and the cosmological constant alone.